

Exercise-1

☒ Marked Questions are Revision Questions.

ONLY ONE OPTION CORRECT TYPE

SECTION - A # SEXUAL REPRODUCTION: INTRODUCTION


1. "Flower is a modified shoot" according to-
 (1) Theophrastus (2) Pliny (3) Goethe (4) Dioscorides
2. Functions of sepals in a flower are -
 (1) Photosynthesis (2) Protection (3) Both (1) and (2) (4) Sporogenesis
- 3.☒ The plants which flower only once in their life-
 (1) Monocarpic (2) Polycarpic (3) Amphicarpic (4) None

SECTION - B # MALE REPRODUCTIVE PART

1. The total nuclei in mature male gametophyte of an angiosperm are
 (1) 2 (2) 3 (3) 4 (4) 5
2. Compound pollengrains are found in
 (1) *Calotropis* (2) *Cyperus* (3) *Typha* (4) None
- 3.☒ Pollinium can be seen in
 (1) *Calotropis* (2) *Coelogynae* (3) *Asclepias* (4) All the above
- 4.☒ Sculpturing on the surface of pollen grain is due to the activity of
 (1) Foot layer and tectum (2) Tectum
 (3) Tectum and sporopollenin (4) Footlayer and Baculate layer.
5. Hay fever (Allergy) is caused due to pollen grains of
 (1) *Amaranthus* (2) *Sorghum* (3) *Ambrosia* (4) All the above.
6. Chromosome number in pollen grain is 6. What shall be it's number in leaf tip cells.
 (1) 6 (2) 12 (3) 24 (4) 3
- 7.☒ There is an abundant occurrence of fossilised pollen grains since it is resistant due to-
 (1) Lignocellulose (2) Sporopollenin (3) Pectocellulose (4) Pectolignin
8. How many pollen mother cells will form 1000 pollen grains?
 (1) 200 (2) 250 (3) 300 (4) 100
9. Monotheous anthers present in-
 (1) Malvaceae (2) Leguminosae (3) Solanaceae (4) Compositae
- 10.☒ Dimorphic tapetum is present in-
 (1) *Typha* (2) *Portulaca*
 (3) *Alectra thomsonii* (4) *Poa*

Reproduction In Flowering Plants

11. In a pollen grain, larger nucleus is-
- (1) Generative nucleus (2) Vegetative nucleus
 (3) Polar nucleus (4) none of these
12. ✘ If sporangia are developed from a single initial cell, the development of sporangia is designated as
- (1) Eusporangiate (2) Leptosporangiate
 (3) Monosporangiate (4) Monocarpic
13. Endothecium, middle layer and tapetum in anther are derived from-
- (1) Primary sporogenous layer (2) Primary parietal layer
 (3) Both (4) None of the above
14. ✘ 'Callase' enzyme which dissolve callose of tetrad of microspores to separate 4 microspores is provided by -
- (1) Pollen grains (2) Middle layer
 (3) Tapetum (4) Endothecium
15. All the cells of anther are diploid except
- (1) Endothelial cells (3) Microspore mother cells
 (2) Epidermal cells (4) Pollen grains
16. Anther of *Arceuthobium* plant is
- (1) Tetra sporangiate (2) Bisporangiate
 (3) Monosporangiate (4) Above (1) and (2) both
17. Linear pollen tetrad is found in
- (1) *Butomopsis* (2) *Polygonum*
 (3) *Magnolia* (4) *Halophila*
18. ✘ Sporopollenin provides resistance to the pollen grain it is chemically
- (1) Protein (2) Fatty substance
 (3) Hetropolysaccharide (4) Homopolysaccharide
19. ✘ #



In the above diagrams identify i, ii, iii and iv and select the suitable options.

- (1) i - Vacuole; ii - symmetrical spindle, iii - Vegetative cell; iv - generative cell
 (2) i - Cytoplasm; ii - Asymmetrical spindle, iii - Generative cell; iv - Vegetative cell
 (3) i - Nuclei; ii - Symmetrical spindle, iii - Tube cell; iv - Vegetative cell
 (4) i - Vacuole; ii - Asymmetrical spindle, iii - Vegetative cell; iv - generative cell

20. Choose incorrect statement.
- (1) In western countries, a large number of pollen products in the form of tablets and syrups are available in market.
 - (2) Some cereals such as rice and wheat, pollen grains lost viability within one year of their release.
 - (3) It is possible to store pollen grains of a large number of species for year in liquid nitrogen (-196°C).
 - (4) Store pollen grains can be used in crop breeding programme.
21. Ubish bodies found in tapetal cells help in formation of
- (1) Pollenkit and pollinia
 - (2) Exine
 - (3) Sporopollenin
 - (4) Intine and pollenkit
22. Largest pollen grain is found in-
- (1) *Halophila*
 - (2) *Myosotis*
 - (3) *Mirabilis*
 - (4) *Lodoicea*
23. Number of prothialial cells in male gametophyte of Angiospermic plant is
- (1) 0
 - (2) 2
 - (3) 3
 - (4) 1

SECTION - C # FEMALE REPRODUCTIVE PART

1. Which of the following is diploid
- (1) Egg
 - (2) Synergids
 - (3) Antipodal cells
 - (4) Secondary nucleus
2. An orthotropous ovule is one, in which micropyle and chalaza are
- (1) At right angles to funicle
 - (2) Parallel to the funicle
 - (3) In straight line of funicle
 - (4) Parallel along with ovule
3. The normal or polygonum type of embryo sac is
- (1) Monosporic and eight nucleate
 - (2) Tetrasporic and six nucleate
 - (3) Monosporic and four nucleate
 - (4) Bisporic and eight nucleate
4. The function of endothelium is
- (1) It protects ovule from toxic substances
 - (2) It helps in fertilization
 - (3) It provides nutrition to embryosac
 - (4) It takes part in deshiscence of Anther
5. Placental or funicular outgrowth present at the micropylar end that directs the passage of pollen tube in to the ovule is
- (1) Aril
 - (2) Caruncle
 - (3) Obturator
 - (4) Raphe
6. A root cell of an angiospermic plant has $2n = 24$ chromosomes. What will no of chromosomes in nucellus cell?
- (1) 12
 - (2) 36
 - (3) 24
 - (4) 18
7. Ovule turns at more than 360° angle is due to excessive growth of funicle in
- (1) Campylotropous ovule
 - (2) Anatropous ovule
 - (3) Orthotropous ovule
 - (4) Circinotropous ovule
8. In which of the following plant, the number of ovules in an ovary may be more than one in
- (1) Wheat
 - (2) Paddy
 - (3) Papaya
 - (4) Mango
9. Polar nuclei are located in-
- (1) Pollen tube
 - (2) Embryo sac
 - (3) Ovule
 - (4) Thalamus
10. The ovule of *capsella* is-

Reproduction In Flowering Plants

- (1) Bitegmic (2) Unitegmic (3) Ategmic (4) Polytegmic

11. In the embryo sac of *Oenothera* no. of antipodal cells are-

- (1) Three (2) One (3) Two (4) None


12. Caruncle is derived from-

- (1) Peduncle (2) Cotyledon (3) Integument (4) none of these

13. Group of lignified cells above the vascular supply of funiculus, which acts as barrier for the growing embryo sac into the base is called-

- (1) Nucellar beak (2) Epistase (3) Hypostase (4) Perisperm

14.# The given figure shows a typical anatropous ovule. What do A, B, C and D represents.



- (1) A →Hilum, B →Funicle, C →Nucellus, D →Micropyle
 (2) A →Hilum, B →Outer integument, C →Nucellus, D →Micropyle
 (3) A →Hilum, B →Outer integument, C →Embryosac, D →Micropyle
 (4) None of these

15. In Angiosperms, the functional megaspore of a linear tetrad is the -

- (1) First nearest to the micropyle (2) Second from the micropyle
 (3) Third from the micropyle (4) Fourth from the micropyle


16. The megasporangium of the angiosperms on maturation gives rise to-

- (1) Fruit (2) Seed (3) Embryo (4) Cotyledon

17. Bisporic type of embryo sac is found in-

- (1) *Polygonum* (2) *Oenothera* (3) *Adoxa* (4) *Allium*

18.# The given figure shows a mature embryo sac. What do A, B, C and D represents.



- (1) A →Embryosac, B →Synergids, C →Central cell, D →Micropylar end
 (2) A →Central cell, B →Synergids, C→Polar nuclei, D→Micropylar end
 (3) A →Synergids, B →Polar nuclei, C →Central cell, D →Filiform apparatus
 (4) A →Central cell, B →Synergids, C→Polar nuclei, D →Filiform apparatus

19. The site of meiotic division in higher plants is

Reproduction In Flowering Plants

- (1) Vegetative buds (2) Root tip cells (3) Stomatal cells (4) Spore mother cells

20. How many cells are present in the female gametophyte of *Capsella* before fertilization?

- (1) 3 (2) 6 (3) 7 (4) Many

SECTION - D # POLLINATION

1. ✖ Flowers never open in

- (1) Chasmogamy (2) Herkogamy (3) Cleistogamy (4) None

2. ✖ Pollination by lever mechanism is found in

- (1) *Ficus* (2) *Calotropis* (3) *Salvia* (4) *Yucca*

3. Hypohydrophily occurs in

- (1) *Vallisneria* (2) *Elodea* (3) *Alisma* (4) *Hydrilla*

4. ✖ Stigma is always rough and sticky in-

- (1) Entomophilous flowers (2) Anemophilous flowers
(3) Hydrophilous flowers (4) All types of flowers

5. ✖ Fragrant flowers with well developed nectaries are an adaptation for-

- (1) Zoophily (2) Anemophily (3) Entomophily (4) Hydrophily

6. Myrmecophily is pollination by-

- (1) Ants (2) Moths (3) Birds (4) Bats

7. Pollination by snails and slugs is-

- (1) Ornithophily (2) Chiropterophily (3) Entomophily (4) Malacophily

8. Some flowers possess pleasant odour and attractive colours for-

- (1) Entomophily (2) Hydrophily (3) Anemophily (4) All of above

9. ✖ From among the situations given below, choose the one that prevents both autogamy and geitonogamy.

- (1) Monoecious plant bearing unisexual flowers
(2) Dioecious plant bearing only male or female flowers
(3) Monoecious plant with bisexual flowers
(4) Dioecious plant with bisexual flowers

10. Anthesis is

- (1) Opening of flower bud (2) pollen mother cell under going meiosis
(3) Dehiscence of Anther (4) Stigma becomes receptive

SECTION - E # FERTILIZATION AND EMBRYOGENESIS, SEED AND POLYEMBRYONY

1. ✖ How many Nucleus participate in double fertilization of *Capsella*

- (1) 2 (2) 5 (3) 3 (4) 4

2. ✖ Development of fruit with out fertilization is

- (1) Parthenocarp (2) Parthenogenesis (3) Sporogamy (4) Autogamy

3. Zygote of *Capsella bursapastoris* divides through

Reproduction In Flowering Plants

- (1) Longitudinal division (2) Equal transverse division
(3) Unequal transverse division (4) Oblique division
4. How many meiosis are required for the formation of 100 grains of wheat?
(1) 100 (2) 200 (3) 150 (4) 125
5. Double fertilization was discovered by Nawaschin in
(1) *Polygonum, Magnolia* (2) *Lilium, Polygonum*
(3) *Fritillaria, Lilium* (4) *Fritillaria, Pepromea*
6. Mature endosperm with irregularity and unevenness in its surface is called Ruminant endosperm. It is found in
(1) Betalnut (2) Maize (3) Coconut (4) Date palm
7. *Casuarina* shows
(1) Porogamy (2) Mesogamy (3) Chalazogamy (4) Acrogamy
8. Which statement is true?
(1) The formation of fruit without fertilization is called parthenocarpy.
(2) The membranous coating of radicle in monocot seed is called coleorhiza.
(3) The development of new individual plant without meiosis and gametic fusion is called Apomixis.
(4) All the above.
9. The phenomenon of pollen tube entering the ovule laterally through integuments is called
(1) Mesogamy (2) Porogamy (3) Chalazogamy (4) None of these
10. Pollen tube enters in embryo sac through
(1) egg cell (2) synergid (3) Antipodal cell (4) Degenerated synergid
11. The effect of pollen grain on colour of endosperm is called
(1) Position effect (2) Warburg effect (3) Metaxenia (4) Xenia
12. Which of the following secrete chemical substances for attracting pollen tube towards micropyle of ovule
(1) Obturator (2) Synergid (3) Filiform apparatus (4) Antipodal cells
13. Syngamy is
(1) Fusion of two cells (2) Fusion of two nuclei
(3) Fusion of two gametes (4) Fusion of two gametic nuclei
14. The fusion product of polar nuclei and male gamete is-
(1) Secondary nucleus (2) Triple fusion
(3) Primary endosperm nucleus (4) Zygote
15. How many meiotic divisions are essential for formation of 100 seeds in cyperaceae family-
(1) 100 (2) 125 (3) 150 (4) 200
16. In angiosperms normally after fertilization

Reproduction In Flowering Plants

- (1) The zygote divides earlier than the primary endosperm nucleus
(2) The primary endosperm nucleus divides earlier than the zygote
(3) Both the zygote and primary endosperm nucleus divide simultaneously
(4) Both the zygote and primary endosperm nucleus undergo a resting period
17. If the number of haploid chromosomes in Gymnosperm is 12, then what will be the number of chromosomes in root and endosperm -
(1) 12, 12 (2) 12, 24 (3) 24, 12 (4) 24, 36
18. In the embryo of a typical dicot and a grass, true homologous structures are:
(1) Coleorhiza and coleoptile (2) Coleoptile and scutellum
(3) Cotyledons and scutellum (4) Hypocotyl and radicle
19. In the flower, if the megaspores forms without meiosis and if one of the megaspores develops into an embryosac, its nuclei would be:
(1) Haploid (2) Diploid
(3) A few haploid and a few diploid (4) With varying ploidy.
20. Which one of the following pairs of plant structure has haploid number of chromosomes?
(1) Nucellus and antipodal cells (2) Egg nucleus and secondary nucleus
(3) Megaspore mother cell and antipodal cell (4) Egg cell and antipodal cell
21. In albuminous seeds the food is stored in
(1) Cotyledons (2) Endosperm (3) Plumule (4) Testa
22. Match the column
- | Column-I | Column-II |
|---------------------------|--|
| (a) Prepotency | (i) shield shaped cotyledon of monocots. |
| (b) Scutellum | (ii) <i>Pinus</i> |
| (c) Translator mechanism | (iii) opening of floral bud |
| (d) Cleavage polyembryony | (iv) <i>Calotropis</i> |
| (e) Anthesis | (v) Apple |
- (1) (a) ii, (b) i, (c) iv, (d) v, (e) iii (2) (a) v, (b) i, (c) iv, (d) iii, (e) ii
(3) (a) iii, (b) v, (c) iv, (d) ii, (e) i (4) (a) v, (b) i, (c) iv, (d) ii, (e) iii
23. Embryology is -
(1) Development of embryo only (2) Mode of gametophyte formation
(3) Sporogenesis and fertilization (4) Sporogenesis, fertilization and embryogenesis
24. If Diploid embryo is directly formed by megaspore mother cell it is called
(1) Non recurrent agamospermy (2) adventive embryony
(3) Diplospory (4) Parthenogenesis
25. Which of the following statement is true.
(1) Pollen tube shows thigmotropic movement before entry in embryo sac.
(2) Perispermic seed is found in castor.
(3) Sporopollenin is proteinaceous substance.
(4) Development of Anther is leptosporangiate type.
26. Polysiphonous pollen tube is a feature of

Reproduction In Flowering Plants

- (1) Cruciferae (2) Asteraceae (3) Cucurbitaceae (4) Liliaceae

27. ✎ Perisperm is

- (1) Outer part of endosperm (2) Destroyed synergid
(3) Destroyed secondary nucleus (4) remain of nucellus

28. ✎ The effect of pollen grain on the outside of endosperm is called.

- (1) Xenia (2) Metaxenia (3) Nemac phenmenon (4) None

29. If the male plant is tetraploid and female plant is diploid. What will be the ploidy level of endosperm after fertilization?

- (1) 3n (2) 4n (3) 5n (4) 6n

30. Which structure develops into seed -

- (1) Ovary (2) Ovule (3) Egg (4) Zygote

MISCELLANEOUS QUESTIONS

1. Which of the following are fleshy fruits.

- (1) Guava, Orange (2) Mango, Mustard (3) Ground nut, Orange (4) All of them

2. How many meiosis are required to produce 50 seeds of tobacco?

- (1) 62 (2) 62.5 (3) 63 (4) 50

3. ✎ A true seed is

- (1) Fertilized ovule (2) Fertilized ovule with embryo
(3) Unfertilized ovule (4) Fertilized ovary

4. Choose correct statements

- (a) Seed typically consists of seed coat(s), cotyledon(s) and embryo axis.
(b) Cotyledons of embryo are simple structures, generally thick and swollen due to storage of food reserve.
(c) Albuminous seeds have no residual endosperm e.g. pea, groundnut.
(d) Micropyle facilitates entry of O_2 and water into the seed during germination.
- (1) a, b (2) b, c (3) a, b, c (4) a, b, d


5. Choose correct statement

- (a) Recent record of 2000 year old viable seed is of the date palm, *Phoenix dactylifera* discovered during the archeological excavation of king herod's palace near dead sea.
(b) Apomixis—special mechanism to produce seed with fertilization.
(c) Flower is modified leaf
(d) Seed have better adaptive strategies for dispersal to new habitats and help the species to colonise in other area.
- (1) a and b (2) a, b, c (3) only a (4) a and d

6. ✎ The position of embryonal axis between plumule and cotyledons is called

- (1) Hypocotyl (2) Epicotyl (3) Coleorhiza (4) Coleoptile

7.#



In the above diagram, identify the correct Labelling and select the correct option

- (1) a - Embryo axis, b - Endosperm, c - Coleorhiza, d - scutellum, e - coleoptile
- (2) a - Radicle, b - Aleuron layer, c - Coleorhiza, d - Endosperm, e - Plumule
- (3) a - Radicle, b - Endosperm, c - Coleorhiza, d - scutellum, e - Plumule
- (4) a - Embryo axis, b - Aleuron layer, c - Root-cap, d - Endosperm, e - Coleoptile

8. Match the column

Column-I

- (a) Helobial endosperm
- (b) Hypophysis
- (c) Ruminant endosperm
- (d) Emasculation
- (e) Mesogamy


Column-II

- (i) *Cucurbita*
- (ii) *Areca*
- (iii) removal of anther from floral bud
- (iv) Radicle
- (v) *Asphodelus*

- (1) a - ii ; b - v ; c - i ; d - iii ; e - iv
- (3) a - i ; b - v ; c - iv ; d - ii ; e - iii

- (2) a - v ; b - iv ; c - ii ; d - iii ; e - i
- (4) a - v ; b - iv ; c - ii ; d - i ; e - iii

9.#



Identify the parts labelled A,B,C,D,E from the list (i- vii) and select the correct options.

Components

- (i) Scutellum
- (ii) Shoot apex
- (iii) Coleoptile
- (iv) Radicle
- (v) Epiblast
- (vi) Coleorhiza
- (vii) Root cap.

- | | A | B | C | D | E |
|-----|-----|-----|-----|----|-----|
| (1) | iii | v | i | vi | vii |
| (2) | iii | vi | vii | ii | iv |
| (3) | ii | vii | vi | i | v |
| (4) | iii | vi | vii | i | iv |

10. The micropyle in a seed helps in the entry of-


Reproduction In Flowering Plants

- (1) Water (2) Pollen tube (3) Male gamete (4) None

11. The tegmen of the seed develops from-

- (1) Perisperm (2) Funicle (3) Inner integument (4) Outer integument

12. The given figure shows false fruits of apple. What do A, B, C and D represents.



- (1) A → Endosperm, B → Thalamus, C → Seed, D → Mesocarp
 (2) A → Thalamus, B → Seed, C → Endocarp, D → Achene
 (3) A → Thalamus, B → Seed, C → Endocarp, D → Mesocarp
 (4) None of these

13. Match column I with column II

Column I

- (A) Albuminous seed
 (B) Non albuminous seed
 (C) Apomixis
 (D) Parthenocarpic fruit

Column II

- (i) Pea, ground nut
 (ii) Wheat, barley
 (iii) Banana
 (iv) Asteraceae and grasses

- (1) A → (i), B → (iv), C → (iii), D → (ii)
 (2) A → (ii), B → (i), C → (iv), D → (iii)
 (3) A → (ii), B → (iv), C → (i), D → (iii)
 (4) A → (ii), B → (iv), C → (iii), D → (i)

14. Choose wrong pair

(1)	Monocarpillary	(i) The gynoecium consists of a single pistil
(2)	Apocarpous	(ii) There are more than one separate pistil
(3)	Funicle	(iii) Stalk by which ovule attach with placenta.
(4)	Scutellum	(iv) It is situated towards both side (Dorsal and ventral of embryo axis)

15. Given below the following statements

- A. Pollen grains are spherical and measures 25 – 50 μm in diameter.
 B. At germ pore sporopollenin is absent
 C. Pollen grain consumption increase performance of athletes and race horses
 D. Pollen grains are shed in two celled stage in more than 60% angiospermic plants

How many statements are wrong –

- (1) 3 (2) 2 (3) 1 (4) 0

16. What will be the ploidy of nucellus, endosperm, female gametophyte and antipodal cells.

- (1) 2n, 3n, n, n (2) n, 3n, 2n, n

Reproduction In Flowering Plants

(3) $2n, 3n, 2n, n$

(4) $3n, 2n, n, n$

17. Mark the incorrect statements-

- A. Dioecy prevents both autogamy and geitonogamy
- B. Self incompatibility prevents inbreeding
- C. Geitonogamy is ecological cross pollination but genetically it is similar to autogamy
- D. Both xenogamy and Geitonogamy decrease inbreeding depression


(1) B

(2) C

(3) A

(4) D

18.# In the given figure A, B, C, D are –



(1) A – epidermis , B – microspore mother cell, C – endothecium, D – tapetum

(2) A – endothecium , B – microspore mother cell, C – middle layer, D – tapetum

(3) A – epidermis , B – microspore mother cell, C – middle layer, D – tapetum

(4) A – endothecium , B – megaspore mother cell, C – endothecium, D – middle layer

19. The correct sequence of embryo formation is-

- (a) heart shaped, globular, mature embryo, proembryo
- (b) proembryo, mature embryo, globular, heart shaped
- (c) globular, proembryo, heart shaped, mature embryo
- (d) proembryo, globular, heart shaped, mature embryo

(1) b

(2) c

(3) a

(4) d

20. How many statements are correct –

- (a) Transfer of pollen grains from the anther to the stigma of another flower of the same plant is known as Geitonogamy.
- (b) Pollination by water is quite rare in flowering plants and is limited to about 30 genera.
- (c) Majority of insect pollinated flowers are large, colourful, fragrant and rich in nectar.
- (d) Pollination guarantee the transfer of the right type of pollen. (Compatible pollen of the same species as the stigma)

(1) One

(2) Two

(3) Three

(4) Four

21. A typical angiospermic anther is

- (1) Bilobed, Monothealous, Tetrasporangiate
- (2) Bilobed, Bisporangiate, Dithecous.
- (3) Bilobed, Dithecous, Tetrasporangiate
- (4) Unilobed, Dithecous, Tetransporangiate

22. All events from pollen deposition on the stigma until pollen tubes enter the ovule are together referred to asA....., which isB..... process. Here A and B are respectively.

(1) A-Embryo sac interaction, B-Dynamic process

Reproduction In Flowering Plants

- (2) A-Pollination, B-Static process
(3) A-pollen pistil interaction, B-static process
(4) A-pollen-pistil interaction, B-Dynamic process
- 23. ✘** Endosperm may either be completely consumed by the developing embryo in plant 'A' before seed maturation or it may persist in mature seed in plant 'B'
A and B plants are respectively.
(1) A-castor, B-pea (2) A-coconut, B-Groundnut
(3) A-pea, B-Groundnut (4) A-Beans, B-castor
- 24. ✘** A typical dicotyledons embryos, consist of an embryonal axis and two cotyledons. The portion of embryonal axis above the level of cotyledons is the 'A' and which terminates with the 'B'
A and B are respectively.
(1) Hypocotyl and Plumule (2) Hypocotyl and Radicle
(3) Epicotyl and Radicle (4) Epicotyl and Plumule
- 25.** A typical microsporangium of angiosperm is surrounded by four wall layers.
I. Epidermis
II. Tapetum
III. Middle layer
IV. Endothecium
The correct sequence from outer to inner side is
(1) II, III, IV, I (2) I, IV, III, II (3) I, II, III, IV (4) IV, III, II, I
- 26. ✘** Perisperm is found in
(1) Beet (2) Black pepper (3) 1 and 2 both (4) In all angiosperms
- 27. ✘** Chasmocleistogamous flowers is/are found in
(a) Viola (common pansy)
(b) Oxalis
(c) Commelina
(1) a only (2) b and c only (3) a and c only (4) a, b, c
- 28. ✘** In _____ of angiosperm pollen grains are shed at 2 celled stage
(1) less than 60% (2) over 60% (3) less than 40% (4) over 90%
- 29.** Anemophilous flowers have
(1) Sessile stigma (2) Small smooth stigma
(3) Coloured flower (4) Large feathery stigma
- 30.** In monocot grafting is almost impossible because they lack
(1) Cambium (2) Ground tissue (3) Vascular bundle (4) Parenchymatous cells
- 31.** The sausage tree (kigelia pinnate) is pollinated by
(1) Bat (2) Bird (3) Wind (4) Water
- 32.** Tapetum is a part of
(1) Male gametophyte (2) Female gametophyte
(3) Ovary wall (4) Anther wall

33. The arrangement of megaspores in a tetrad in an angiosperm is
(1) Decussate (2) Tetrahedral (3) Linear (4) Isobilateral
34. Egg apparatus consists of
(1) Egg (2) Egg and Polar nuclei
(3) Egg and Synergids (4) Egg and Antipodal cells
35. What is the liquid part of green coconut?
(1) Endosperm (2) Female gametophyte
(3) Nucellus (4) Embryo
36. Pollen grains are
(1) Male gamete (2) Male gametophyte
(3) Non-functional megaspore (4) Nucelus
37. Seed coat is formed by
(1) Integument (2) Nucellus (3) fruit wall (4) None
38. Wind pollinated flowers are
(1) Small, scented and colourless (2) Small, nonscented and colourless
(3) Big, scented and coloured (4) Big, nonscented and colourless
39. Virus free culture is got from
(1) Primary root (2) Pith of stem (3) apical cells (4) Lamina cells.
40. Radicle end of embryo is towards
(1) Hilum (2) Chalaza (3) Funicle (4) Micropyle
41. In a monocot, endosperm cells have 24 chromosomes. What shall be the chromosome number in embryo?
(1) 24 (2) 16 (3) 12 (4) 8
42. Ovule integument gets transformed into
(1) Seed (2) Seed coat (3) Fruit wall (4) Cotyledons
43. In 82% of angiosperm families, ovule is
(1) Anatropous (2) Orthotropous (3) Amphitropous (4) Circinotropous
44. Tapetal cells of stamens are
(1) Diploid, uninucleate (2) Tetraploid, binucleate
(3) Hexaploid, tetranucleate (4) Polyploid, multinucleate
45. Vegetative fertilization, which involves formation of endosperm, is fusion of
(1) One male gamete with diploid secondary nucleus
(2) Two vegetative cells
(3) Two male gametes
(4) Female gamete with secondary nucleus.
46. Largest cell of the ovule is
(1) Megaspore mother cell (2) Antipodal cell
(3) Central cell (4) Size of cells variable

Reproduction In Flowering Plants

47. Device for self pollination is
(1) Heterostyly (2) Dicliny (3) Unisexuality (4) None of the above
48. Rarely in angiosperms, the pollen tube develops further in embryo sac. The abnormality is called
(1) Metaxenia (2) Nemec phenomenon
(3) Xenia (4) Mesogamy
49. In flowering plants, meiosis takes place during
(1) Pollen grain formation (2) Seed formation
(3) Gamete formation (4) Seed germination
50. Which of the following statements is true with reference to cross pollination
(1) It most often results in higher yield of plants
(2) It occurs only in unisexual flowers
(3) It can fail to occur due to distance
(4) It requires production of large number of pollen grains
51. Pollen grains are produced in
(1) Ovary (2) Pollen sac (3) Filament (4) Stigma
52. For self pollination, a flower should be
(1) Asexual (2) Monosexual (3) Unisexual (4) Bisexual
53. Part which is grafted on the stump of another plant is called
(1) Graft (2) Bulbil (3) Scion (4) Stock
54. Grafting is not successful in monocots but is successful in dicots because they have
(1) Vascular bundles arranged in a ring
(2) Cambium for secondary growth
(3) Vessels with elements arranged end to end
(4) Cork cambium
55. Match the columns
- | Columns-I | Columns-II |
|--|--|
| a. Zoophily | 1. Pollination by birds |
| b. Ornithophily | 2. Pollination by insects |
| c. Entomophily | 3. Pollination by bats |
| d. Chiropterophily | 4. Pollination by animals. |
| (1) (a) — (3), (b) — (2), (c) — (1), (d) — (4) | (2) (a) — (1), (b) — (2), (c) — (3), (d) — (4) |
| (3) (a) — (4), (b) — (1), (c) — (2), (d) — (3) | (4) (a) — (4), (b) — (2), (c) — (1), (d) — (3) |
56. In angiosperms, endosperm is formed by
(1) Division of fused polar nuclei
(2) Free nuclear division of megaspore
(3) Division of fused synergids and male gamete
(4) Division of fused polar nuclei and male gamete
57. What would be number of chromosomes in aleurone layer if megaspore mother cell contains 10 chromosomes?
(1) 10 (2) 20 (3) 15 (4) None of the above
58. During development of male gametophyte from pollen mother cell, there occurs

Reproduction In Flowering Plants


- (1) Two meiotic divisions and one mitotic division
(2) Two mitotic divisions
(3) One meiotic and two mitotic division
(4) One meiotic cell division and one mitotic cell division
59. Male gametes are formed by
(1) Pollen cell (2) Generative cell (3) Pollen tube cell (4) Pollen mother cell
60. Embryo sac develops from megaspore mother cell through
(1) 1 meiosis and 2 mitosis (2) 1 meiosis and 3 mitosis
(3) 1 meiosis and two mitosis (4) 1 meiosis and 2 mitosis.
61. Versatile anthers are connected with
(1) Entomophily (2) Malacophily (3) Ornithophily (4) Anemophily
62. Eight nucleate embryosacs are
(1) Always monosporic
(2) Always bisporic
(3) Always tetrasporic
(4) Sometimes monosporic, sometimes, bisporic and sometimes tetrasporic.
63. Double fertilization leading to initiation of endosperm in angiosperms requires
(1) Fusion of one polar nucleus and second male gamete only
(2) Fusion of two polar nuclei and second male gamete only
(3) Fusion of 4 or more polar nuclei and second male gamete only
(4) All the above type of fusions in different types of angiosperms.
64. Anemophilous type of pollination is found in
(1) Coconut (2) *Salvia* (3) Bottle brush (4) *Vallisneria*
65. Entry of pollen tube through micropyle is
(1) Chalazogamy (2) mesogamy (3) Porogamy (4) Pseudogamy
66. Plants with poor root system are propagated through
(1) Layering (2) Leaf cutting (3) Stem cutting (4) Grafting
67. Nutritive layer of anther wall is
(1) Epidermis (2) Endothecium (3) tapetum (4) Archesporium
68. Suspensor of embryo is formed by
(1) Basal cell (2) Apical cell (3) Terminal cell (4) Hypophysis
69. In the monocotyledonous seeds the endosperm is separated from the embryo by a distinct layer known as
(1) Testa (2) Tegmen (3) Aleurone layer (4) Scutellum
70. Choose the mismatched option
(1) Wind-*Cannabis*-anemophily (2) Water-*Zostera*-hydrophily
(3) Insects-*Salvia*-entomophily (4) Birds-*Adansonia*-ornithophily
71. The seed which have separate endosperm

Reproduction In Flowering Plants

- (1) maize (2) Onion (3) Rice (4) Bean
72. For production of haploids, we culture
(1) Shoot tip (2) anther (3) root tip (4) None of these
73. What does the filiform apparatus do at the entrance into ovule?
(1) It helps in the entry of pollen tube into a synergid
(2) It prevents entry of more than one pollen tube into the embryo sac
(3) It brings about opening of the pollen tube
(4) It guides pollen tube from a synergid to egg
74. A plant cell has potential to develop into a full plant. This is called
(1) totipotency (2) gene cloning (3) tissue culture (4) regeneration
75. Xenia refers to effect of pollen on
(1) stem (2) taste of fruit (3) vascular tissue (4) endosperm
76. 200 seeds are produced from how many fruits of maize
(1) 200 (2) 100 (3) 50 (4) 5
77. An orthotropous ovule has micropyle and chalaza
(1) parallel to funicle (2) parallel alongwith ovule
(3) at right angle to funicle (4) in straight line with funicle
78. Perisperm is
(1) outer part to embryo sac (2) degenerate synergid
(3) degenerate secondary nucleus (4) remains of nucellus
79. ~~79.~~ Stem cutting is employed in the propagation of-
(1) Banana (2) Mango (3) Sugarcane (4) Cotton
80. ~~80.~~ The most common method of Vegetative propagation described by ancient gardeners is-
(1) Layering (2) Gootee (3) Grafting (4) Ground layering
81. Bulbils occur is-
(1) *Cycas* (2) *Agave* (3) *Dioscorea* (4) All the above
82. ~~82.~~ In a grafted plant, stock has 48 chromosomes while scion has 24 chromosomes. The chromosome number for root cells and eggs are-
(1) 48 and 24 (2) 24 and 24 (3) 24 and 12 (4) 48 and 12
83. In vegetative propagation by tubers, which of following remains constant through generations-
(1) Morphology (2) Vigour only
(3) Vigour and morphology only (4) Morphology, vigour and disease resistance

Exercise-2

1. Which of the following characteristics of a flower would attract humming birds for pollination but not bees **(NSEB -2010)**
 (i) Fragrant flowers (ii) Great amount of nectar
 (iii) Long tubular flowers (iv) Deep-seated nectary
 (v) Petals forming a lip for resting (vi) Yellow petals
 (1) (i), (ii) and (v) only (2) (i), (iii) and (iv) only (3) (ii), (iv) and (vi) only (4) (i) and (ii) only
2. Filiform apparatus found during development in angiosperms is a thickening on the **(NSEB -2010)**
 (1) antipodals (2) polar nuclei (3) egg (4) synergids
3. Which of the following is the correct combination of merits of an inflorescence? **(NSEB -2011)**
 i. Flowers can be unisexual
 ii. Increased efficiency of pollination
 iii. Ensuring self pollination and fertility
 iv. Attract pollinators easily
 (1) i,ii and iv (2) ii,iii and iv (3) ii and iii (4) ii and iv
4. Arrange the embryo-sac development stages of angiosperms in correct order: **(INBO - 2012)**



- (1) v → i → iv → ii → iii → vii → vi → viii
 (2) viii → v → ii → iv → iii → vii → vi → i
 (3) i → ii → iv → v → viii → iii → vii → vi
 (4) viii → i → v → ii → iv → iii → vi → vii

Exercise-3

PART - I : NEET / AIPMT QUESTION (PREVIOUS YEARS)

1. Which is the characteristics for Ornithophily
 (1) Scented flowers
 (2) Bright red coloured flowers and inflorescence
 (3) White coloured funnel shaped large corolla
 (4) Yellow flower with nectaries at the base of the corolla tube

Reproduction In Flowering Plants

2. Double fertilization involves
(1) Fusion of secondary nucleus with both male gametes.
(2) Fusion of secondary nucleus with one male gamete.
(3) Fusion of one nucleus with one male gamete.
(4) Any of the above
3. Anemophily occurs in
(1) *Salvia* (2) Bottle brush (3) Date palm (4) *Vallisneria*
4. What is the direction of micropyle in anatropous ovule
(1) Upward (2) Downward (3) Right (4) left
5. Which type of association is found in between entomophilous flower and pollinating agent
(1) Mutualism (2) Commensalism (3) Co-operation (4) Co-evolution
6. In flowering plants Archegonium gives rise to
(1) Wall of sporangium (2) Both wall and sporogenous cells
(3) Wall and tapetum (4) Tapetum and sporogenous cells
7. A diploid female plant is crossed with tetraploid male. The ploidy of endosperm will be
(1) Tetraploidy (2) Pentaploidy (3) Triploidy (4) Diploidy
8. Secondary nucleus present in the middle of embryo sac is
(1) Tetraploid (2) Triploid (3) Diploid (4) Haploid
9. In oogamy, fertilization occurs between
(1) Small non-motile female gamete and large motile male gamete
(2) Large non-motile female gamete and small motile male gamete
(3) A large non-motile female gamete and a small non-motile male gamete
(4) A large motile female gamete and a small non-motile male gamete
10. In which one part both the plants can be vegetatively propagated by leaf segments
(1) *Agave* and *Kalanchoe* (2) *Bryophyllum* and *Kalanchoe*
(3) *Asparagus* and *Bryophyllum* (4) *Chrysanthemum* and *Agave*
11. In a type of apomixis known as adventitious embryony, embryos develop directly from
(1) Nucellus or integument (2) Zygote
(3) Synergids of antipodals of embryo sac (4) Accessory embryo sac in the ovule.
12. Arrangement of nuclei in normal dicot embryo sac is
(1) 3 + 3 + 2 (2) 2 + 4 + 2 (3) 3 + 2 + 3 (4) 3 + 3 + 3
13. What would be number of chromosomes in the cells of aleurone layer in plant species have 8 chromosomes in its synergids
(1) 8 (2) 16 (3) 24 (4) 32.

Reproduction In Flowering Plants

14. Parthenocarpic fruits are produced by
(1) Treating plants with phenyl Mercuric acetate
(2) Treating plants with low concentrations of gibberellic acid and auxin
(3) Removing androecium of flowers before release of pollen grains
(4) Raising plants from vernalised seeds.
15. Which one is surrounded by callose wall
(1) Male gamete (2) Pollen grain
(3) Egg (4) Microspore mother cell.
16. Endosperm is consumed by developing embryo in the seed of
(1) Coconut (2) Pea (3) Maize (4) Castor
17. Unisexuality of flowers prevents
(1) Geitonogamy but not xenogamy (2) Autogamy and geitonogamy
(3) Autogamy but not geitonogamy (4) Both geitonogamy and xenogamy
18. Which one of the following is resistant to enzyme action
(1) Pollen exine (2) Leaf cuticle (3) Cork (4) Wood fibre
19. An example of a seed with endosperm, perisperm, and caruncle is:
(1) castor (2) cotton (3) coffee (4) lily
20. Apomictic embryos in *citrus* arise from
(1) Maternal sporophytic tissue in ovule (2) Antipodal cells
(3) Diploid egg (4) Synergids
21. Wind pollinated flowers are
(1) Small, producing large number of dry pollen grains
(2) Large producing abundant nectar and pollen
(3) Small, producing nectar and dry pollen
(4) Small, brightly coloured, producing large number of pollen grains
22. Transfer of pollen grains from the anther to the stigma of another flower of the same plant is called
(1) Geitonogamy (2) Karyogamy (3) Autogamy (4) Xenogamy
23. 'Filiform' apparatus is a characteristic feature of:
(1) Suspensor (2) Egg (3) Synergid (4) Zygote
24. Nucellar polyembryony is reported in species
(1) *Citrus* (2) *Gossypium* (3) *Triticum* (4) *Brassica*
25. In which one of the following pollination is autogamous
(1) Geitonogamy (2) Xenogamy (3) Chasmogamy (4) Cleistogamy
26. Wind pollination is common in:
(1) Legumes (2) Lilies (3) Grasses (4) Orchids

Reproduction In Flowering Plants

27. Both, autogamy and geitonogamy are prevented in -
(1) Papaya (2) Cucumber (3) Castor (4) Maize
28. An organic substance that can withstand environmental extremes and cannot be degraded by any enzyme is :
(1) Cuticle (2) Sporopollenin (3) Lignin (4) Cellulose
29. The gynoecium consists of many free pistils in flowers of
(1) *Aloe* (2) Tomato (3) *Papaver* (4) *Michelia*
30. Even in absence of pollinating agents seed setting is assured in
(1) *Commellina* (2) *Zostera* (3) *Salvia* (4) Fig
31. Which one of the following statements is wrong?
(1) When pollen is shed at two-celled stage, double fertilization does not take place.
(2) Vegetative cell is larger than generative cell.
(3) Pollen grains in some plants remain viable for months.
(4) Intine is made up of cellulose and pectin.
32. Plants with ovaries having only one or a few ovules, are generally pollinated by:
(1) Bees (2) Butterflies (3) Birds (4) Wind
33. What is the function of germ pore?
(1) Emergence of radicle
(2) Absorption of water for seed germination
(3) Initiation of pollen tube
(4) Release of male gametes
34. Perisperm differs from endosperm in
(1) having no reserve food
(2) being a diploid tissue
(3) its formation by fusion of secondary nucleus with several sperms
(4) being a haploid tissue
35. Megasporangium is equivalent to:
(1) Fruit (2) Nucellus (3) Ovule (4) Embryo sac
36. Advantage of cleistogamy is :
(1) More vigorous offspring (2) No dependence of pollinators
(3) Vivipary (4) Higher genetic variability
37. Which one of the following statements is correct?
(1) Sporogenous tissue is haploid (2) Endothecium produces the microspores
(3) Tapetum nourishes the developing pollen (4) Hard outer layer of pollen is called intine
38. Which one of the following statements is correct?
(1) The seed in grasses is not endospermic.
(2) Mango is a parthenocarpic fruit
(3) A proteinaceous aleurone layer is present in maize grain.
(4) A sterile pistil is called a staminode.

Reproduction In Flowering Plants

39. Geitonogamy involves
- (1) Fertilization of a flower by the pollen from another flower of the same plant
 - (2) Fertilization of a flower by the pollen from same flower.
 - (3) Fertilization of a flower by the pollen from a flower of another plant in the same population
 - (4) Fertilization of a flower by the pollen from a flower of another plant belonging to a distant population
40. Male gametophyte with least number of cells is present in:
- (1) *Pteris*
 - (2) *Funaria*
 - (3) *Lilium*
 - (4) *Pinus*
41. Pollen tablets are available in the market for:
- (1) In vitro fertilization
 - (2) Breeding programmes
 - (3) Supplementing food
 - (4) Ex situ conservation
42. Function of filiform apparatus is to:
- (1) Recognize the suitable pollen at stigma
 - (2) Stimulate division of generative cell
 - (3) Producer nectar
 - (4) Guide the entry of pollen tube
43. Non- albuminous seed is produced in:
- (1) Maize
 - (2) Castor
 - (3) Wheat
 - (4) Pea
44. Which one of the following statements is not true?
- (1) Pollen grains of some plants cause severe allergies and bronchial afflictions in some people
 - (2) The flowers pollinated by flies and bats secrete foul odour to attract them
 - (3) Honey is made by bees by digesting pollen collected from flowers
 - (4) Pollen grains are rich in nutrients, and they are used in the form of tablets and syrups
45. The hilum is a scar on the:
- (1) Fruit, where it was attached to pedicel
 - (2) Fruit, where style was present
 - (3) Seed, where micropyle was present
 - (4) Seed, where funicle was attached
46. Which one of the following may require pollinators, but is genetically similar to autogamy?
- (1) Xenogamy
 - (2) Apogamy
 - (3) Cleistogamy
 - (4) Geitonogamy
47. Which of the following are the important floral rewards to the animal pollinators?
- (1) Nectar and pollen grains
 - (2) Floral fragrance and calcium crystals
 - (3) Protein pellicle and stigmatic exudates
 - (4) Colour and large size of flower
48. Transmission tissue is characteristic feature of
- (1) Solid style
 - (2) Dry stigma
 - (3) Wet stigma
 - (4) Hollow style
49. Male gametophyte in angiosperms produces:
- (1) Single sperm and a vegetative cell
 - (2) Single sperm and two vegetative cells
 - (3) Three sperms
 - (4) Two sperms and a vegetative cell
50. Coconut water from a tender coconut is:
- (1) Free nuclear endosperm
 - (2) Innermost layers of the seed coat
 - (3) Degenerated nucellus
 - (4) Immature embryo

Reproduction In Flowering Plants

51. Flowers are unisexual in:
(1) Cucumber (2) China rose (3) Onion (4) Pea
52. Which one of the following fruits is parthenocarpic?
(1) Apple (2) Jackfruit (3) Banana (4) Brinjal
53. Filiform apparatus is characteristic feature of:
(1) Nucellar embryo (2) Aleurone cell (3) Synergids (4) Generative cell
54. The wheat grain has an embryo with one, large, shield-shaped cotyledon known as:
(1) Coleorrhiza (2) Scutellum (3) Coleoptile (4) Epiblast
55. In angiosperms, microsporogenesis and megasporogenesis:
(1) form gametes without further divisions (2) Involve meiosis
(3) occur in ovule (4) occur in anther
56. Proximal end of the filament of stamen is attached to the:
(1) Thalamus or petal (2) Anther
(3) Connective (4) Placenta
57. The coconut water from tender coconut represents:
(1) Free nuclear endosperm (2) Endocarp
(3) Fleshy mesocarp (4) Free nuclear proembryo
58. Which of the following statements is not correct?
(1) Some reptiles have also been reported as pollinators in some plant species. ,
(2) Pollen grains of many species can germinate on the stigma of a flower, but only one pollen tube of the same species grows into the style.
(3) Insects that consume pollen or nectar without bringing about pollination are called pollen / nectar robbers.
(4) Pollen germination and pollen tube growth are regulated by chemical components of pollen interacting with those of the pistil
59. Seed formation without fertilization in flowering plants involves the process of:
(1) Apomixis (2) Sporulation (3) Budding (4) Somatic hybridization
60. Which one of the following statements is **not** true?
(1) Stored pollen in liquid nitrogen can be used in the crop breeding programmes
(2) Tapetum helps in the dehiscence of anther
(3) Exine of pollen grains is made up of sporopollenin
(4) Pollen grains of many species cause severe allergies
61. Which one of the following in and generates new genetic combinations leading to variation?
(1) Nucellar polyembryony (2) Vegetative reproduction
(3) Parthenogenesis (4) Sexual reproduction

62. Match **Column-I** with **Column-II** and select the correct option using the codes given below:

Column-I

- a. Pistils fused together
- b. Formation of gametes
- c. Hyphae of higher Ascomycetes
- d. Unisexual female flower

Column-II

- (i) Gametogenesis
- (ii) Pistillate
- (iii) Syncarpous
- (iv) Dikaryotic

Codes:

	a	b	c	d
(1)	(iii)	(i)	(iv)	(ii)
(2)	(iv)	(iii)	(i)	(ii)
(3)	(ii)	(i)	(iv)	(iii)
(4)	(i)	(ii)	(iv)	(iii)

63. In majority of angiosperms

- (1) a small central cell is present in the embryo sac:
- (2) egg has a filiform apparatus
- (3) there are numerous antipodal cells
- (4) reduction division occurs in the megaspore mother cells

64. Pollination in water hyacinth and water lily is brought about by the agency of

- (1) bats
- (2) water
- (3) insects or wind
- (4) birds

65. The ovule of an angiosperm is technically equivalent to

- (1) megaspore
- (2) megasporangium
- (3) megasporophyll
- (4) megaspore mother cell

66. Flowers which have single ovule in the ovary and are packed into inflorescence are usually pollinated by:

- (1) Water
- (2) Bee
- (3) Wind
- (4) Bat

67. A dioecious flowering plant prevents both:

- (1) Autogamy and xenogamy
- (2) Autogamy and geitonogamy
- (3) Geitonogamy and xenogamy
- (4) Cleistogamy and xenogamy

68. Attractants and rewards are required for:

- (1) Anemophily
- (2) Entomophily
- (3) Hydrophily
- (4) Cleistogamy

69. Winged pollen grains are present in

- (1) Mustard
- (2) Pinus
- (3) Mango
- (4) Cycas

70. Which of the following has proved helpful in preserving pollen as fossils?

- (1) Pollenkitt
- (2) Sporopollenin
- (3) Oil content
- (4) Cellulosic intine

71. Persistent nucellus in the seed is known as

- (1) Tegmen
- (2) Chalaza
- (3) Perisperm
- (4) Hilum

72. In some plants, the female gamete develops into embryo without fertilization. This phenomenon is known as:

- (1) Parthenogenesis
- (2) Autogamy
- (3) Parthenocarpy
- (4) Syngamy

73. What is the fate of the male gametes discharged in the synergid?

- (1) One fuses with the egg and other fuses with central cell nuclei.
- (2) One fuses with the egg, other(s) degenerate in the synergid.
- (3) All fuse with the egg.
- (4) One fuses with the egg, other (s) fuse (s) with synergid nucleus.

Reproduction In Flowering Plants

74. Which one of the following statements regarding post-fertilization development in flowering plants is incorrect ?
(1) Ovules develop into embryo sac (2) Ovary develops into fruit
(3) Zygote develops into embryo (4) Central cell develops into endosperm
75. Which is the most common type of embryo sac in angiosperms?
(1) Tetrasporic with one mitotic stage of divisions
(2) Monosporic with three sequential mitotic divisions
(3) Monosporic with two sequential mitotic divisions
(4) Bisporic with two sequential mitotic divisions
76. What type of pollination takes place in Vallisneria ?
(1) Pollination occurs in submerged condition by water.
(2) Flowers emerge above surface of Water and pollination occurs by insects.
(3) Flowers emerge above water surface and pollen is carried by wind.
(4) Male flowers are carried by water currents to female flowers at surface of water.
77. In which one of the following, both autogamy and geitonogamy are prevented?
(1) Wheat (2) Papaya (3) Castor (4) Maize


PART - II : AIIMS QUESTION (PREVIOUS YEARS)

1. The cotyledon in monocot plant is one and
(1) Lateral (2) Terminal (3) Basal (4) Vertical
2. The root cell of wheat plant has 42 chromosomes. What would be the number of chromosomes in the synergid cells
(1) 7 (2) 14 (3) 21 (4) 28
3. Pollen grain are able to tolerate extremes of temperature and desiccation because their exine consists of
(1) Cutin (2) Suberin (3) Sporopollenin (4) Callose
4. Pollen tube usually enters the embryo sac through
(1) one of the synergids (2) Directly penetrates the egg
(3) Between one synergid and central cell (4) By knocking of antipodal cells
5. Double fertilization involves
(1) Fertilization of egg by two male gametes
(2) Fertilization of two eggs in the same embryo sac by two sperms brought by one pollen tube
(3) Fertilization of the egg and the central cell by two sperms brought by the same pollen tube
(4) Fertilization of the egg and the central cell by two sperms brought by two different pollen tubes.
6. A scion is grafted on a stock. Quality of fruits produced will depend upon genotypes of
(1) Scion (2) Stock (3) 1 and 2 both (4) None of the above
7. Match the following ovular structure with post fertilization structure and select the correct alternative.
(a) Ovule 1. Endosperm
(b) Funiculus 2. Aril
(c) Nucellus 3. Seed
(d) Polar nuclei 4. Perisperm
(1) a-3, b-2, c-4, d-1 (2) a-3, b-2, c-1, d-4
(3) a-1, b-2, c-3, d-4 (4) a-2, b-3, c-1, d-4

Reproduction In Flowering Plants

8. The plant part which consists of two generations one within the other, is
(1) Embryo (2) Germinated pollen grain
(3) Unfertilized ovule (4) Seed
9. Which one of the following pairs of plant structures has haploid number of chromosomes?
(1) nucellus and antipodal cells. (2) egg nucleus and secondary nucleus.
(3) megaspore mother cell and antipodal cells. (4) egg cell and antipodal cells.

- 10.# The given figure shows L.S of the seed of maize. What do A, B, C and D represent?



- (1) A : endosperm, B : scutellum, C : plumule, D : coleoptile
(2) A : scutellum, B : pericarp, C : radicle, D : coleoptile
(3) A : endosperm, B : scutellum, C : radicle, D : coleorrhiza
(4) A : scutellum, B : peri carp, C : plumule, D : coleorrhiza
11. The normal type of embryo sac is 8-nucleated and
(1) 8-celled (2) 7-celled (3) 6-celled (4) 5-celled
12. Embryo sac is
(1) megaspore (2) microgametophyte
(3) female gametophyte (4) megasporangium
13. Feathery stigma belongs to
(1) Wheat (2) Pea (3) *Datura* (4) *Caesalpinia*
14. A fertilized egg of a plant has 40 chromosomes. The number of chromosomes present in the microspore mother cell is
(1) 20 (2) 40 (3) 60 (4) 80
15. Chasmogamy refers to the condition where
(1) Flowers remains closed (2) Flowers are absent
(3) Flowers are open (4) Flower are gamopetalous
16. What is common between vegetative reproduction and apomixis?
(1) Both are applicable to only dicot plants
(2) Both bypass the flowering phase
(3) Both occur around the year
(4) Both produce progeny identical to the parent
17. Emasculation is not required when flowers are
(1) bisexual (2) intersexual (3) unisexual (4) either (a) or (b)
18. How many haploid nuclei are present in a mature pollen grain?
(1) One (2) Two (3) Three (4) Four

Answers

EXERCISE - 1

SECTION - A

1. (3) 2. (3) 3. (1)

SECTION - B

1. (2) 2. (3) 3. (4) 4. (2) 5. (4) 6. (2) 7. (2)
 8. (2) 9. (1) 10. (3) 11. (2) 12. (2) 13. (2) 14. (3)
 15. (4) 16. (3) 17. (4) 18. (2) 19. (4) 20. (2) 21. (2)
 22. (3) 23. (1)

SECTION - C

1. (4) 2. (3) 3. (1) 4. (3) 5. (3) 6. (3) 7. (4)
 8. (3) 9. (2) 10. (1) 11. (4) 12. (3) 13. (3) 14. (1)
 15. (4) 16. (2) 17. (4) 18. (4) 19. (4) 20. (3)

SECTION - D

1. (3) 2. (3) 3. (4) 4. (1) 5. (3) 6. (1) 7. (4)
 8. (1) 9. (2) 10. (1)

SECTION - E

1. (2) 2. (1) 3. (3) 4. (4) 5. (3) 6. (1) 7. (3)
 8. (4) 9. (1) 10. (4) 11. (4) 12. (3) 13. (3) 14. (3)
 15. (4) 16. (2) 17. (3) 18. (3) 19. (2) 20. (4) 21. (2)
 22. (4) 23. (4) 24. (3) 25. (2) 26. (3) 27. (4) 28. (2)
 29. (2) 30. (2)

MISCELLANEOUS QUESTIONS

1. (1) 2. (3) 3. (2) 4. (4) 5. (4) 6. (2) 7. (3)
 8. (2) 9. (4) 10. (1) 11. (3) 12. (3) 13. (2) 14. (4)
 15. (4) 16. (1) 17. (4) 18. (2) 19. (4) 20. (3) 21. (3)
 22. (4) 23. (4) 24. (4) 25. (2) 26. (3) 27. (4) 28. (2)
 29. (4) 30. (1) 31. (1) 32. (4) 33. (3) 34. (3) 35. (1)
 36. (2) 37. (1) 38. (2) 39. (3) 40. (4) 41. (2) 42. (2)
 43. (1) 44. (4) 45. (1) 46. (3) 47. (4) 48. (2) 49. (1)
 50. (4) 51. (2) 52. (4) 53. (3) 54. (2) 55. (3) 56. (4)
 57. (3) 58. (3) 59. (2) 60. (2) 61. (4) 62. (4) 63. (2)
 64. (1) 65. (3) 66. (4) 67. (3) 68. (1) 69. (3) 70. (4)
 71. (1) 72. (2) 73. (1) 74. (1) 75. (4) 76. (1) 77. (4)
 78. (4) 79. (3) 80. (3) 81. (4) 82. (4) 83. (4)

EXERCISE - 2

1. (3) 2. (4) 3. (4) 4. (4)

EXERCISE - 3

PART - I

1. (4) 2. (2) 3. (3) 4. (2) 5. (1) 6. (2) 7. (1)
 8. (3) 9. (2) 10. (2) 11. (1) 12. (3) 13. (3) 14. (2)
 15. (4) 16. (2) 17. (3) 18. (1) 19. (1) 20. (1) 21. (1)
 22. (1) 23. (3) 24. (1) 25. (4) 26. (3) 27. (1) 28. (2)
 29. (4) 30. (1) 31. (1) 32. (4) 33. (3) 34. (2) 35. (3)
 36. (2) 37. (3) 38. (3) 39. (1) 40. (3) 41. (3) 42. (4)
 43. (4) 44. (3) 45. (4) 46. (4) 47. (1) 48. (1) 49. (4)
 50. (1) 51. (1) 52. (3) 53. (3) 54. (2) 55. (2) 56. (1)
 57. (1) 58. (2) 59. (1) 60. (2) 61. (4) 62. (1) 63. (4)
 64. (3) 65. (2) 66. (3) 67. (2) 68. (2) 69. (2) 70. (2)
 71. (3) 72. (1) 73. (1) 74. (1) 75. (2) 76. (4) 77. (2)

PART - II

1. (1) 2. (3) 3. (3) 4. (1) 5. (3) 6. (1) 7. (1)
 8. (4) 9. (4) 10. (3) 11. (2) 12. (3) 13. (1) 14. (2)
 15. (3) 16. (4) 17. (3) 18. (2) 19. (2) 20. (3) 21. (1)
 22. (3) 23. (4) 24. (4)